

INDUCED SUBGRAPHS AND TREE DECOMPOSITIONS

VII. BASIC OBSTRUCTIONS IN H -FREE GRAPHS

TARA ABRISHAMI^{*†}, BOGDAN ALECU^{**¶}, MARIA CHUDNOVSKY^{**‡}, SEPEHR HAJEBI[§],
AND SOPHIE SPIRKL^{§||}

ABSTRACT. We say a class \mathcal{C} of graphs is *clean* if for every positive integer t there exists a positive integer $w(t)$ such that every graph in \mathcal{C} with treewidth more than $w(t)$ contains an induced subgraph isomorphic to one of the following: the complete graph K_t , the complete bipartite graph $K_{t,t}$, a subdivision of the $(t \times t)$ -wall or the line graph of a subdivision of the $(t \times t)$ -wall. In this paper, we adapt a method due to Lozin and Razgon (building on earlier ideas of Weißauer) to prove that the class of all H -free graphs (that is, graphs with no induced subgraph isomorphic to a fixed graph H) is clean if and only if H is a forest whose components are subdivided stars.

Their method is readily applied to yield the above characterization. However, our main result is much stronger: for every forest H as above, we show that forbidding certain connected graphs containing H as an induced subgraph (rather than H itself) is enough to obtain a clean class of graphs. Along the proof of the latter strengthening, we build on a result of Davies and produce, for every positive integer η , a complete description of unavoidable connected induced subgraphs of a connected graph G containing η vertices from a suitably large given set of vertices in G . This is of independent interest, and will be used in subsequent papers in this series.

Keywords. Induced subgraph, Tree decomposition, Treewidth.

1. INTRODUCTION

Background. All graphs in this paper are finite and simple.

Treewidth is a well-studied graph parameter that is of great interest in both structural and algorithmic graph theory. It was notably featured in the seminal work of Robertson and Seymour on graph minors [18], and in numerous other papers ever since. For a more in-depth overview of the literature, the reader is invited to see, for example, Bodlaender’s survey [9] and the references therein.

As a part of their graph minors series, Robertson and Seymour fully described the unavoidable minors in graphs of large treewidth. The relevant result, the so-called Grid Theorem [19], states that every graph of large enough treewidth must contain a minor isomorphic to a large grid, or equivalently, a subgraph isomorphic to a large wall (the $(t \times t)$ -wall, denoted by $W_{t \times t}$, is a planar graph of maximum degree three on $2t^2 - 2t$ vertices; see [2] for a precise definition and

^{*}PRINCETON UNIVERSITY, PRINCETON, NJ, USA

^{**}SCHOOL OF COMPUTING, UNIVERSITY OF LEEDS, LEEDS, UK

[§]DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA

[†] SUPPORTED BY NSF-EPSRC GRANT DMS-2120644.

[‡] SUPPORTED BY NSF-EPSRC GRANT DMS-2120644 AND BY AFOSR GRANT FA9550-22-1-0083.

[¶] SUPPORTED BY DMS-EPSRC GRANT EP/V002813/1.

^{||} WE ACKNOWLEDGE THE SUPPORT OF THE NATURAL SCIENCES AND ENGINEERING RESEARCH COUNCIL OF CANADA (NSERC), [FUNDING REFERENCE NUMBER RGPIN-2020-03912]. CETTE RECHERCHE A ÉTÉ FINANCÉE PAR LE CONSEIL DE RECHERCHES EN SCIENCES NATURELLES ET EN GÉNIE DU CANADA (CRSNG), [NUMÉRO DE RÉFÉRENCE RGPIN-2020-03912]. THIS PROJECT WAS FUNDED IN PART BY THE GOVERNMENT OF ONTARIO.

Date: November 7, 2023.

THIS IS AN ACCEPTED MANUSCRIPT. THE PUBLISHED VERSION APPEARED IN THE JOURNAL OF COMBINATORIAL THEORY, SERIES B VOLUME 164, JANUARY 2024, PAGES 443–472 AND IS AVAILABLE HERE: [HTTPS://DOI.ORG/10.1016/J.JCTB.2023.10.008](https://doi.org/10.1016/j.jctb.2023.10.008)




FIGURE 1. The 3-basic obstructions, including a subdivision of $W_{3 \times 3}$ (middle) and its line graph (right).

see Figure 1). Since walls have large treewidth themselves, and treewidth cannot increase when taking minors, that result gives a structural dichotomy: a graph has large treewidth if and only if it contains a large wall as a subgraph.

The overarching goal of the current series and of several other recent works [1, 7, 15, 16, 20, 21] is to understand treewidth from the perspective of induced subgraphs rather than minors. A first remark is that to force bounded treewidth, we need to forbid four kinds of induced subgraphs: a complete graph K_t , a complete bipartite graph $K_{t,t}$, all subdivisions of the $(t \times t)$ -wall $W_{t \times t}$ for some t , and the line graphs of all subdivisions of $W_{t \times t}$ for some t . Let us call these graphs the t -basic obstructions (see Figure 1), and say that a graph G is t -clean if G contains no induced subgraph isomorphic to a t -basic obstruction. Moreover, we say a class \mathcal{C} of graphs is *clean* if the treewidth of t -clean graphs in \mathcal{C} is bounded from above by a function of t .

The class of all graphs is not clean: various constructions of unbounded treewidth avoiding the basic obstructions have been discovered [7, 10, 20]. In fact, it is at the moment unclear whether a dichotomy similar to the Grid Theorem is at all achievable for induced subgraphs. Nevertheless, steady progress is being made. Of note is the following result, characterizing all finite sets of graphs which yield bounded treewidth when forbidden as induced subgraphs:

Theorem 1.1 (Lozin and Razgon [16]). *Let \mathcal{H} be a finite set of graphs. Then the class of all graphs with no induced subgraph isomorphic to a member of \mathcal{H} has bounded treewidth if and only if \mathcal{H} contains a complete graph, a complete bipartite graph, a forest of maximum degree at most three in which every component has at most one vertex of degree more than two, and the line graph of such a forest.*

In addition, several clean classes have been identified. For instance, Aboulker, Adler, Kim, Sintiari and Trotignon [1] proved that every proper minor-closed class of graphs is clean:

Theorem 1.2 (Aboulker, Adler, Kim, Sintiari and Trotignon [1]). *For every graph H , the class of all graphs with no minor isomorphic to H is clean. Equivalently, for every graph H and integers $t \geq 1$, there exists an integer $\xi = \xi(H, t) \geq 1$ such that every graph with no minor isomorphic to H and treewidth more than ξ contains either a subdivision of $W_{t \times t}$ or the line graph of a subdivision of $W_{t \times t}$ as an induced subgraph.*

They also conjectured that graph classes of bounded maximum degree are clean, which was later proved by Korhonen [15]:

Theorem 1.3 (Korhonen [15]). *For every integer $d \geq 1$, the class of graphs of maximum degree at most d is clean. Equivalently, for all integers $d, t \geq 1$, there exists an integer $\gamma = \gamma(d, t) \geq 1$ such that every graph with maximum degree at most d and treewidth more than γ contains either a subdivision of $W_{t \times t}$ or the line graph of a subdivision of $W_{t \times t}$ as an induced subgraph.*

There are also a number of results concerning holes, where a *hole* in a graph is an induced cycle of length at least four. In particular, it was shown that (even hole, diamond, pyramid)-free graphs are clean [3], and graphs in which no vertex has two or more neighbors in a hole disjoint from itself are clean [4]. It was also independently proved twice that graphs with no long hole




FIGURE 2. The smallest tree that is not a subdivided star.

are clean. For every positive integer λ , let \mathcal{H}_λ be the class of all graphs with no hole of length more than λ .

Theorem 1.4 (Gartland, Lokshtanov, Pilipczuk, Pilipczuk and Rzażewski [13], Weißauer [21]). *For every integer $\lambda \geq 1$, the class \mathcal{H}_λ is clean.*

Our results. The main result of this paper is Theorem 4.1. The precise statement of Theorem 4.1 requires some set-up, and we postpone it until Section 4. Informally, we show that every t -clean graph of sufficiently large treewidth contains, as an induced subgraph, a “connec-tification” of any given subdivided star forest F . Roughly speaking, this is a graph which can be partitioned into a “rooted” copy of F and a second part, which only attaches at the roots of F and “minimally connects” these roots.

The proof of Theorem 4.1 uses three ingredients. The first one is Theorem 8.1, which adapts the methods from [16] (itself employing the strategy from [21]) in order to show that clean graphs with a large *block* – a certain kind of highly connected structure – must contain a large subdivided star forest. As a byproduct of this, we also obtain another way to derive Theorem 1.4.

The second ingredient is Theorem 6.5. This theorem combines a result of Weißauer linking blocks and tree decompositions, together with Korhonen’s bounded degree result (Theorem 1.3), in order to show that the class of graphs without a large block is clean.

The final ingredient, Theorem 5.2, is a result of independent interest, and will be used in future papers in our series. Starting from a result of Davies [11], we provide a complete description of minimal connected graphs containing many vertices from a suitably large subset of a connected component. Put differently, we show that if a large enough set of vertices belongs to the same component, then a large subset of them are contained in one of a few prescribed induced subgraphs.

We note that the first two out of those intermediate results already yield (the difficult direction of) an appealing dichotomy for clean classes defined by one forbidden induced subgraph. Indeed, writing \mathcal{F}_H for the class of graphs with no induced subgraph isomorphic to H , we prove:

Theorem 1.5. *Let H be a graph. Then \mathcal{F}_H is clean if and only if H is a subdivided star forest.*

While the stronger Theorem 4.1 might appear unwieldy at first, we remark that it has easier-to-state implications that are still more general than the above dichotomy. To illustrate this, denote by $\tilde{\mathcal{F}}_H$ the class of all graphs with no induced subgraph isomorphic to a subdivision of H . It follows that the “if” direction of Theorem 1.5 is equivalent to $\tilde{\mathcal{F}}_H$ being clean for every subdivided star forest H , and Theorem 1.4 is equivalent to $\tilde{\mathcal{F}}_H$ being clean for every cycle H . Then Theorem 4.1 readily implies the following, where by a *subdivided double star*, we mean a tree with at most two vertices of degree more than two.

Theorem 1.6. *Let H be a forest in which one component is a subdivided double star and every other component is a subdivided star. Then $\tilde{\mathcal{F}}_H$ is clean.*

We remark that a full grid-type theorem for induced subgraphs is equivalent to a characterization of families \mathcal{H} of graphs for which the class of all \mathcal{H} -free graphs is clean. This remains out of reach, and Theorem 1.5 takes the first step towards answering this question by characterizing all singletons \mathcal{H} for which the class of all \mathcal{H} -free graphs is clean.

Here is a natural next step: *for which finite families \mathcal{H} of graphs is the class of all \mathcal{H} -free graphs clean?* From Theorem 1.5, it follows that such a finite set \mathcal{H} containing a subdivided star

forest has the above property. One may then speculate that in fact *all* finite set of graphs with the above property must contain a subdivided star forest. This, however, is false: for instance, assume that H is the unique double star on six vertices (see Figure 2; note that H is the smallest tree that is not a subdivided star). Then $\mathcal{H} = \{H, K_3\}$ has the above property; in fact, this follows from the main result of an upcoming paper [6] where the last four authors of the present work provide a full description of finite families \mathcal{H} for which the class of all \mathcal{H} -free graphs is clean.

Outline of the paper. We set up our notation and terminology in Section 2. Section 3 describes the construction of [10], which is used to prove the “only if” direction of Theorem 1.5. In Section 4, we state Theorem 4.1 precisely, and show how to deduce Theorems 1.5 and 1.6 from it. In Section 5, we show that a connected graph G with a sufficiently large subset S of its vertices contains an induced connectifier with many vertices from S . The main result of Section 6 is Theorem 6.5, where we prove that the class of graphs with no k -block is clean. In Section 7, we show that in a t -clean graph, every huge block can be transformed into a large block such that there is no short path between any two vertices of the new block. Section 8 uses this in order to show that a t -clean graph with a huge block contains a large subdivided star forest. Finally, in Section 9, we combine the main results from Sections 5, 6 and 8 to prove Theorem 4.1.

2. PRELIMINARIES

Graphs, subgraphs, and induced subgraphs. All graphs in this paper are finite and with no loops or multiple edges. Let $G = (V(G), E(G))$ be a graph. A *subgraph* of G is a graph obtained from G by removing vertices or edges, and an *induced subgraph* of G is a graph obtained from G by only removing vertices. Given a subset $X \subseteq V(G)$, $G[X]$ denotes the subgraph of G induced by X , that is, the graph obtained from G by removing the vertices not in X . We put $G \setminus X = G[V(G) \setminus X]$ (and in general, we will abuse notation and use induced subgraphs and their vertex sets interchangeably). Additionally, for an edge $e \in E(G)$, we write $G - e$ to denote the graph obtained from G by removing the edge e . For a graph H , by a *copy of H in G* , we mean an induced subgraph of G isomorphic to H , and we say G *contains H* if G contains a copy of H . We also say G is *H -free* if G does not contain H . For a class \mathcal{H} of graphs we say G is \mathcal{H} -free if G is H -free for every $H \in \mathcal{H}$. For a graph H , we write $G = H$ whenever G and H have the same vertex set and the same edge set.

Neighborhoods. Let $v \in V(G)$. The *open neighborhood* of v , denoted by $N(v)$, is the set of all vertices in G adjacent to v . The *closed neighborhood* of v , denoted by $N[v]$, is $N(v) \cup \{v\}$. Let $X \subseteq G$. The *open neighborhood* of X , denoted by $N(X)$, is the set of all vertices in $G \setminus X$ with at least one neighbor in X . If H is an induced subgraph of G and $X \subseteq G$ with $H \cap X = \emptyset$, then $N_H(X) = N(X) \cap H$. Let $X, Y \subseteq V(G)$ be disjoint. We say X is *complete* to Y if all possible edges with one end in X and one end in Y are present in G , and X is *anticomplete* to Y if there is no edge between X and Y . In the case $X = \{x\}$, we often say x is *complete* (*anticomplete*) to Y to mean X is complete (anticomplete) to Y .

Tree decompositions and blocks. A *tree decomposition* (T, χ) of G consists of a tree T and a map $\chi : V(T) \rightarrow 2^{V(G)}$ with the following properties:

- (i) For every vertex $v \in V(G)$, there exists $t \in V(T)$ such that $v \in \chi(t)$.
- (ii) For every edge $v_1v_2 \in E(G)$, there exists $t \in V(T)$ such that $v_1, v_2 \in \chi(t)$.
- (iii) For every vertex $v \in V(G)$, the subgraph of T induced by $\{t \in V(T) \mid v \in \chi(t)\}$ is connected.

For each $t \in V(T)$, we refer to $\chi(t)$ as a *bag* of (T, χ) . The *width* of a tree decomposition (T, χ) , denoted by $\text{width}(T, \chi)$, is $\max_{t \in V(T)} |\chi(t)| - 1$. The *treewidth* of G , denoted by $\text{tw}(G)$, is the minimum width of a tree decomposition of G .

Cliques, stable sets, paths, and cycles. A *clique* in G is a set of pairwise adjacent vertices in G , and a *stable set* in G is a set of pairwise non-adjacent vertices in G . A *path in G* is an induced subgraph of G that is a path, while a *cycle in G* is a (not necessarily induced) subgraph of G that is a cycle. If P is a path, we write $P = p_1 \cdots p_k$ to mean that $V(P) = \{p_1, \dots, p_k\}$, and p_i is adjacent to p_j if and only if $|i - j| = 1$. We call the vertices p_1 and p_k the *ends of P* , and say that P is *from p_1 to p_k* . The *interior of P* , denoted by P^* , is the set $P \setminus \{p_1, p_k\}$. For a path P in G and $x, y \in P$, we denote by $P[x, y]$ the subpath of P with ends x and y . The *length* of a path P is the number of its edges. Let C be a cycle. We write $C = c_1 \cdots c_k c_1$ to mean $V(C) = \{c_1, \dots, c_k\}$, and c_i is adjacent to c_j if and only if $|i - j| \in \{1, k - 1\}$. A *hole of G* is an induced subgraph of G that is a cycle. The *length* of a cycle or a hole is the number of its edges.

Subdivisions. By a *subdivision* of a graph G , we mean a graph obtained from G by replacing the edges of G by pairwise internally disjoint paths between the corresponding ends. Let $r \geq 0$ be an integer. An *r -subdivision* of G is a subdivision of G in which the path replacing each edge has length $r + 1$. Also, a *$(\leq r)$ -subdivision* of G is a subdivision of G in which the path replacing each edge has length at most $r + 1$, and a *$(\geq r)$ -subdivision* of G is defined similarly. We refer to a (≥ 1) -subdivision of G as a *proper* subdivision of G .

Classes of graphs. A class \mathcal{C} of graphs is called *hereditary* if it is closed under isomorphism and taking induced subgraphs, or equivalently, if \mathcal{C} is the class of all \mathcal{H} -free graphs for some family \mathcal{H} of graphs. For a class of graphs \mathcal{C} and a positive integer t , we denote by \mathcal{C}^t the class of all t -clean graphs in \mathcal{C} . Thus, \mathcal{C} is clean if for every positive integer t there exists a positive integer $w(t)$ such that every graph in \mathcal{C}^t has treewidth at most $w(t)$. The following is immediate from the definition of a clean class.


Lemma 2.1. *Let \mathcal{X} be a class of graphs. Assume that for every t , there exists a clean class of graphs \mathcal{Y}_t such that $\mathcal{X}^t \subseteq \mathcal{Y}_t$. Then \mathcal{X} is clean. In particular, every subclass of a clean class is clean.*

Forests and stars. By a *branch vertex* of a graph G , we mean a vertex of degree more than two in G . For every forest F , we say a vertex $v \in V(F)$ is a *leaf* of F if v has degree at most one in F . We denote by $\mathcal{L}(F)$ the set of all leaves of F . By a *star* we mean a graph isomorphic to the complete bipartite graph $K_{1,\delta}$ for some integer $\delta \geq 0$, and a *star forest* is a forest in which every component is a star. Then subdivided stars are exactly trees with at most one branch vertex, and subdivided star forests are exactly forests in which every component is a subdivided star. A *subdivided double star* is a tree with at most two branch vertices.

By a *rooted subdivided star S* we mean a subdivided star S together with a choice of one vertex r in S , called the *root*, such that if S is not a path, then r is the unique branch vertex of S . A *rooted subdivided star forest F* is a subdivided star forest with a choice of a root for every component of F . We also refer to the root of each component of F as a *root of F* , and denote by $\mathcal{R}(F)$ the set of all roots of F . By a *stem* in F , we mean a path in F from a leaf to a root. It follows that each stem is the (unique) path from a leaf of some component of F to the root of the same component. The *reach* of a rooted subdivided star S is the maximum length of a stem in S . Also, the *reach* of a subdivided star forest F is the maximum reach of its components and the *size* of F is the number of its components. For a positive integer θ and graph H , we denote by θH the disjoint union of θ copies of H . For integers $\delta \geq 0$ and $\lambda \geq 1$, we denote by $S_{\delta,\lambda}$ the $(\lambda - 1)$ -subdivision of $K_{1,\delta}$. So for $\delta \geq 3$, $\theta S_{\delta,\lambda}$ is a subdivided star forest of maximum degree δ , reach λ and size θ .

3. A CONSTRUCTION FROM [10]

The goal of this section is to prove the “only if” direction of Theorem 1.5 using a construction from [10].

FIGURE 3. The graphs $J_{0,1,4}$ (left) and $J_{1,1,4}$ (right).

We begin with a definition, which will be used in subsequent sections, as well. Let P be a path and $\rho, \sigma \geq 0$ and $\theta \geq 1$ be integers. A 2θ -tuple $(p_1, \dots, p_{2\theta})$ of vertices of P is said to be a (ρ, σ) -widening of P if

- the vertices p_1 and $p_{2\theta}$ are the ends of P ;
- traversing P from p_1 to $p_{2\theta}$, the vertices $p_1, \dots, p_{2\theta}$ appear on P in this order;
- $P[p_{2i-1}, p_{2i}]$ has length ρ for each $i \in [\theta]$, and;
- $P[p_{2i}, p_{2i+1}]$ has length at least σ for each $i \in [\theta - 1]$.

The (ρ, σ) -widening $(p_1, \dots, p_{2\theta})$ is *strict* if for each $i \in [\theta - 1]$, $P[p_{2i}, p_{2i+1}]$ has length equal to σ . Also, we say a θ -tuple (p_1, \dots, p_θ) of vertices of P is a σ -widening of P if the 2θ -tuple $(p_1, p_1, \dots, p_\theta, p_\theta)$ is a $(0, \sigma)$ -widening of P .

We now describe the construction of [10] (though [10] only mentions the case $\rho = 0$). Let $\rho \geq 0$, $\sigma \geq 1$ and $\theta \geq 2$ be integers. We define $J = J_{\rho, \sigma, \theta}$ to be the graph with the following specifications (see Figure 3).

- J contains θ pairwise disjoint and anticomplete paths P_1, \dots, P_θ .
- For each $j \in [\theta]$, P_j admits a strict (ρ, σ) -widening $(p_1^j, \dots, p_{2\theta}^j)$.
- We have $J \setminus (\bigcup_{i \in [\theta]} V(P_i)) = \{x_1, \dots, x_\theta\}$ such that x_1, \dots, x_θ are all distinct, and for all $i, j \in [\theta]$, we have $N_J(x_i) = \bigcup_{j \in [\theta]} P_j[p_{2i-1}^j, p_{2i}^j]$.

The following was proved in [11]. Here we include a proof for the sake of completeness.

Theorem 3.1. *For all integers $\rho \geq 0$, $\sigma \geq 1$ and $\theta \geq 2$, $J_{\rho, \sigma, \theta}$ is a 4-clean graph of treewidth at least θ .*

Proof. Note that $J_{\rho, \sigma, \theta}$ contains a $K_{\theta, \theta}$ -minor (by contracting each path P_i into a vertex), which implies that $\text{tw}(J_{\rho, \sigma, \theta}) \geq \theta$. Also, $J_{\rho, \sigma, \theta}$ is easily seen to be $\{K_4, K_{3,3}\}$ -free. Let us say that a connected graph H is *feeble* if either H has a vertex v such that $H \setminus N_H[v]$ is not connected, or H has a set S of at most two branch vertices such that $H \setminus S$ has maximum degree at most two. Then every connected induced subgraph of $J_{\rho, \sigma, \theta}$ is feeble. On the other hand, for an integer $t \geq 4$, let H be either a subdivision of $W_{t \times t}$ or the line graph of such a subdivision. Then one may observe that for every vertex $v \in H$, $H \setminus N_H[v]$ is connected. Moreover, H contains a stable set S of branch vertices with $|S| \geq 3$. It follows that H is not feeble, and so H is not isomorphic to an induced subgraph of $J_{\rho, \sigma, \theta}$. Hence, $J_{\rho, \sigma, \theta}$ is 4-clean, as desired. ■

The proof of the next lemma is straightforward, and we leave it to the reader.

Lemma 3.2. *For all integers $\sigma \geq 1$ and $\theta \geq 2$, the following hold.*

- $J_{0, \sigma, \theta}$ has girth at least $2\sigma + 4$.
- Let $u_1, u_2 \in J_{1, \sigma, \theta}$ such that for each $i \in \{1, 2\}$, $N_{J_{1, \sigma, \theta}}(u_i)$ contains a stable set of cardinality three. Then there is no path of length less than $\sigma + 2$ in $J_{1, \sigma, \theta}$ from u_1 to u_2 .

We are now ready to prove the main result of this section.

Theorem 3.3. *Let H be a graph for which \mathcal{F}_H is clean. Then H is a subdivided star forest.*

Proof. By the assumption, for every integer $t \geq 1$, there exists an integer $w(t) \geq 1$ such that every t -clean graph in \mathcal{F}_H has treewidth at most $w(t)$. We deduce:

(1) H is a forest.

Suppose not. Let σ be the length of the shortest cycle in H . By Theorem 3.1, $J_{0,\sigma,w(4)+1}$ is 4-clean. Also, by the first outcome of Lemma 3.2, $J_{0,\sigma,w(4)+1}$ has girth at least $2\sigma + 4$, and so $J_{0,\sigma,w(4)+1} \in \mathcal{F}_H$. But then we have $\text{tw}(J_{0,\sigma,w(4)+1}) \leq w(4)$, which violates Theorem 3.1. This proves (1).

(2) Every component of H has at most one branch vertex.

Suppose for a contradiction that some component C of H contains two branch vertices u and v . By (1), H is a forest, and so C is a tree. Therefore, there exists a unique path in H from u to v , say of length σ , and we have $|N_H(u) \setminus N_H(v)|, |N_H(v) \setminus N_H(u)| \geq 2$. It follows from the second outcome of Lemma 3.2 that $J_{1,\sigma,w(4)+1} \in \mathcal{F}_H$. Also, by Theorem 3.1, $J_{1,\sigma,w(4)+1}$ is 4-clean. But then we have $\text{tw}(J_{1,\sigma,w(4)+1}) \leq w(4)$, a contradiction with Theorem 3.1. This proves (2).

Now the result follows from (1) and (2). This completes the proof of Theorem 3.3. \blacksquare

4. CONNECTIFICATION AND STATEMENT OF THE MAIN RESULT

Here we state the main result of the paper, Theorem 4.1. Then we discuss how it implies Theorems 1.5 and 1.6.

We need numerous definitions. A vertex v of a graph G is said to be *simplicial* if $N_G(v)$ is a clique of G . The set of all simplicial vertices of G is denoted by $\mathcal{Z}(G)$. It follows that every degree-one vertex in G belongs to $\mathcal{Z}(G)$. In particular, for every forest F , we have $\mathcal{L}(F) = \mathcal{Z}(F)$.

By a *caterpillar* we mean a tree C of maximum degree three in which all branch vertices lie on a path. A path P in C is called a *spine* for C if all branch vertices of C belong to $V(P)$ and subject to this property P is maximal with respect to inclusion (our definition of a caterpillar is non-standard for two reasons: a caterpillar is often allowed to be of arbitrary maximum degree, and a spine often contains all vertices of degree more than one.)

Let C be a caterpillar with $\theta \geq 3$ leaves. Note that C has exactly $\theta - 2$ branch vertices, and both ends of each spine of C are leaves of C . Also, for every leaf $l \in \mathcal{L}(C)$, there exists a unique branch vertex in C , denoted by v_l , for which the unique path in C from l to v_l does not contain any branch vertex of C other than v_l (and, in fact, $\{v_l : l \in \mathcal{L}(C)\}$ is the set of all branch vertices of C). We say an enumeration (l_1, \dots, l_θ) of $\mathcal{L}(C) = \mathcal{Z}(C)$ is σ -wide if for some spine P of C , the θ -tuple $(l_1, v_{l_2}, \dots, v_{l_{\theta-1}}, l_\theta)$ is a σ -widening of P . Also, let H be the line graph of C . Then assuming e_l to be the unique edge in C incident with the leaf $l \in \mathcal{L}(C)$, we have $\mathcal{Z}(H) = \{e_l : l \in \mathcal{L}(C)\}$. An enumeration $(e_{l_1}, \dots, e_{l_\theta})$ of $\mathcal{Z}(H)$ is called σ -wide if (l_1, \dots, l_θ) is a σ -wide enumeration of $\mathcal{L}(C)$. By a σ -caterpillar, we mean a caterpillar C for which $\mathcal{L}(C)$ admits a σ -wide enumeration. It follows that if H is the line graph of a caterpillar C , then $\mathcal{Z}(H)$ admits a σ -wide enumeration if and only if C is a σ -caterpillar.

Let H be a graph and S be a set. We say H is S -tied if $\mathcal{Z}(H) \subseteq H \cap S$ and *loosely S -tied* if $\mathcal{Z}(H) = H \cap S$. Also, for a positive integer $\eta \geq 1$, we say H is (loosely) (S, η) -tied if H is (loosely) S -tied and $|H \cap S| = \eta$. It follows that if H is loosely (S, η) -tied, then $|\mathcal{Z}(H)| = \eta$.

For a graph G , a set $S \subseteq G$ and integers $\eta \geq 2$ and $\sigma \geq 1$ and $i \in \{0, \dots, 4\}$, we say an induced subgraph H of G is an (S, η, σ) -connectifier of type i if H satisfies the condition (Ci) below.

- (C0) H is a loosely (S, η) -tied line graph of a subdivided star in which every stem has length at least σ .
- (C1) H is an (S, η) -tied rooted subdivided star with root r in which every stem has length at least σ , and we have $(H \cap S) \setminus \mathcal{L}(H) \subseteq \{r\}$.
- (C2) H is an (S, η) -tied path with $H \cap S = \{s_1, \dots, s_\eta\}$ where (s_1, \dots, s_η) is a σ -widening of H .
- (C3) H is a loosely (S, η) -tied σ -caterpillar.




FIGURE 4. From left to right: an $(S, 4)$ -connectifier H of type 0, 1, 2, 3 and 4. Circled nodes depict the vertices in $H \cap S$. Note that for the subdivided star, r may or may not belong to S (and if it does, then we have $\eta = 5$).

(C4) H is a loosely (S, η) -tied line graph of a σ -caterpillar.

See Figure 4. We say H is an (S, η) -connectifier of type i if it is an $(S, \eta, 1)$ -connectifier of type i . Also, we say H is an (S, η, σ) -connectifier (resp. (S, η) -connectifier) if it is an (S, η, σ) -connectifier (resp. (S, η) -connectifier) of type i for some $i \in \{0, \dots, 4\}$.

Note that connectifiers of type 0 contain large cliques, and since we mostly work with t -clean graphs, they do not come up in our arguments. However, for the sake of generality, we cover them in both the above definition and the main result of the next section, Theorem 5.2. We also remark that, unlike the connectifiers of other types, connectifiers of type 1 in fact need to be “tied” rather than “loosely tied.” For instance, let G be a subdivided star with root r and let $S = \mathcal{L}(G) \cup \{r\}$. Then for every $\eta > 1$, every (S, η) -connectifier in G contains r .

Let σ be a positive integer, F be a graph and $X \subseteq F$ with $|X| \geq 2$. Let $\pi : [|X|] \rightarrow X$ be a bijection. By a σ -connectification of (F, X) with respect to π , we mean a graph Ξ with the following specifications.

- F is an induced subgraph of Ξ .
- $F \setminus X$ is anticomplete to $\Xi \setminus F$.
- Let $H = \Xi \setminus (V(F) \setminus X)$. Then H is $(X, |X|, \sigma)$ -connectifier in Ξ of type i for $i \in [4]$ such that
 - if H is of type 2 (that is, H is path), then, traversing H from one end to another, $(\pi(1), \dots, \pi(|X|))$ is a σ -widening of H , and;
 - if H is of type 3 or 4, then $(\pi(1), \dots, \pi(|X|))$ is a σ -wide enumeration of $\mathcal{Z}(H)$.

Also, by a σ -connectification of (F, X) , we mean a σ -connectification of (F, X) with respect to some bijection $\pi : [|X|] \rightarrow X$.

Let $\mathcal{C}_{\sigma, F, X, \pi}$ be the class of all graphs with no induced subgraph isomorphic to a σ -connectification of (F, X) with respect to π , and $\mathcal{C}_{\sigma, F, X}$ be the class of all graphs with no induced subgraph isomorphic to a σ -connectification of (F, X) . In other words, $\mathcal{C}_{\sigma, F, X}$ is the intersection of all classes $\mathcal{C}_{\sigma, F, X, \pi}$ over all bijections $\pi : [|X|] \rightarrow X$. As a result, for every $\pi : [|X|] \rightarrow X$, we have $\mathcal{C}_{\sigma, F, X} \subseteq \mathcal{C}_{\sigma, F, X, \pi}$.

The following is our main result, which we will prove in Section 9.

Theorem 4.1. *Let $\sigma \geq 1$ be an integer, F be a rooted subdivided star forest of size at least two and $\pi : [|R(F)|] \rightarrow R(F)$ be a bijection. Then the class $\mathcal{C}_{\sigma, F, R(F), \pi}$ is clean.*

Next we discuss briefly how to deduce Theorems 1.5 and 1.6 using Theorem 4.1. The “only if” direction of Theorem 1.5 is proved in Theorem 3.3. Also, the “if” direction of Theorem 1.5 follows from Theorem 1.6. So it suffices to prove Theorem 1.6, which we restate:

Theorem 4.2. *Let H be a forest in which one component is a subdivided double star and every other component is a subdivided star. Then $\tilde{\mathcal{F}}_H$ is clean.*

Proof. We define F and σ as follows. If H is a subdivided star forest, then let $F = 2H$ be rooted and $\sigma = 2$. If H is not a subdivided star forest, let H' be the 1-subdivision of H . Then there are two branch vertices $u_1, u_2 \in H'$ and a path Q in H' from u_1 to u_2 with $Q^* \neq \emptyset$ such that $F' = H' \setminus Q^*$ is a subdivided star forest. For each $i \in \{1, 2\}$, let F_i be the component of F'

containing u_i . Then u_i is a vertex of maximum degree in F_i and so u_i is a valid choice for a root of F_i . Let F' be rooted such that $u_1, u_2 \in \mathcal{R}(F')$. Let δ, λ and θ be the maximum degree, the reach and the size of F' , respectively. So we have $\delta, \theta \geq 2$ and $\lambda \geq 1$. Let $F = \theta S_{\delta+1, \lambda}$ be rooted with its unique choice of roots and let $\sigma = |Q| \geq 3$. Then, every σ -connectification of $(F, \mathcal{R}(F))$ contains a subdivision of H . Therefore, for every bijection $\pi : [|\mathcal{R}(F)|] \rightarrow \mathcal{R}(F)$, we have $\tilde{\mathcal{F}}_H \subseteq \mathcal{C}_{\sigma, F, \mathcal{R}(F)} \subseteq \mathcal{C}_{\sigma, F, \mathcal{R}(F), \pi}$. It follows that for every integer $t \geq 1$, we have $\tilde{\mathcal{F}}_H^t \subseteq \mathcal{C}_{\sigma, F, \mathcal{R}(F), \pi}$. This, together with Theorem 4.1 and Lemma 2.1, implies Theorem 4.2. ■

In fact, one may deduce Theorem 1.5 directly using the material from Sections 3, 6, 7 and 8 (and in particular, skipping Section 5).

5. OBTAINING A CONNECTIFIER

We begin with the following folklore result, see, for example, [2] for a proof.

Theorem 5.1. *Let G be a connected graph, $X \subseteq V(G)$ with $|X| = 3$ and H be a connected induced subgraph of G with $X \subseteq H$ and with H minimal subject to inclusion. Then one of the following holds.*

- *There exists a vertex $a \in H$ and three paths $\{P_x : x \in X\}$ (possibly of length zero) where P_x has ends a and x , such that*
 - $H = \bigcup_{x \in X} P_x$, and;
 - the sets $\{P_x \setminus \{a\} : x \in X\}$ are pairwise disjoint and anticomplete.
- *There exists a triangle with vertex set $\{a_x : x \in X\}$ in H and three paths $\{P_x : x \in X\}$ (possibly of length zero) where P_x has ends a_x and x , such that*
 - $H = \bigcup_{x \in X} P_x$;
 - the sets $\{P_x \setminus \{a_x\} : x \in X\}$ are pairwise disjoint and anticomplete, and;
 - for distinct $x, y \in X$, $a_x a_y$ is the only edge of H between P_x and P_y .

Theorem 5.1 may be reformulated as follows: for every choice of three vertices x, y, z in a connected graph G , there is an induced subgraph H of G containing x, y, z such that, for some $\delta \in [3]$, H is isomorphic to either a subdivision of $K_{1, \delta}$ or the line graph of a subdivision of $K_{1, \delta}$, and $\mathcal{Z}(H) \subseteq \{x, y, z\}$. The main result of this section, the following, can be viewed as a qualitative extension of Theorem 5.1.

Theorem 5.2. *For every integer $\eta \geq 1$, there exists an integer $\mu = \mu(\eta) \geq 1$ with the following property. Let G be a graph and $S \subseteq V(G)$ with $|S| \geq \mu$ such that S is contained in a connected component of G . Then G contains an (S, η) -connectifier H . In particular, H is connected, $|H \cap S| = \eta$, and every vertex in $H \cap S$ has degree at most η in H .*

For a graph G , $S \subseteq G$ and positive integer η , one may observe that (S, η) -connectifiers are minimal with respect to being connected and containing η vertices from S . Also, for $\eta_1, \eta_2 \geq 4$ (which, given Theorem 5.1, captures the main content of Theorem 5.2) and distinct $i_1, i_2 \in \{0, 1, \dots, 4\}$, no (S, η_1) -connectifier of type i_1 contains an induced subgraph which is an (S, η_2) -connectifier of type i_2 . Therefore, Theorem 5.2 provides an efficient characterization of all minimally connected induced subgraphs of G containing many vertices from a sufficiently large subset S of vertices in G .

In order to prove Theorem 5.2, we need a few definitions and a result from [11]. By a *big clique* in a graph J , we mean a maximal clique of cardinality at least three. A graph J is said to be a *bloated tree* if

- every edge of J is contained in at most one big clique of J .
- for every big clique K of J and every $v \in K$, v has at most one neighbor in $J \setminus K$; and
- the graph obtained from J by contracting each big clique into a vertex is a tree.

It follows that every bloated tree is connected, and every connected induced subgraph of a bloated tree is a bloated tree. Furthermore, we deduce:

Lemma 5.3. *Let J be a bloated tree. Then for every cycle C in J , $V(C)$ is a clique of J .*

Proof. Suppose for a contradiction that for some cycle C in J , $V(C)$ contains two vertices which are non-adjacent in J . Let C be chosen with $|V(C)| = k$ as small as possible. It follows that $k \geq 4$. Let $C = c_1 \cdots c_k c_1$ such that c_1 and c_i are not adjacent for some $i \in \{3, \dots, k-1\}$. Let P be a path in J from c_1 to c_i with $P^* \subseteq \{c_2, \dots, c_{i-1}\}$ and let Q be a path in J from c_1 to c_i with $Q^* \subseteq \{c_{i+1}, \dots, c_k\}$. So P and Q are internally vertex-disjoint and $|P|, |Q| \geq 3$. Also, $H = J[P \cup Q]$ is a connected induced subgraph of J , and so H is a bloated tree. If P^* is anticomplete to Q^* , then H is cycle. But then the graph obtained from H by contracting each big clique into a vertex is H itself, which is not tree, a contradiction with H being a bloated tree. It follows that there exists $p \in P^*$ and $q \in Q^*$ such that $pq \in E(J)$. Consequently, $C_1 = c_1 - P - p - q - Q - c_1$ and $C_2 = c_i - P - p - q - Q - c_i$ are two cycles in J with $|V(C_1)|, |V(C_2)| < |V(C)|$. Thus, by the choice of C , for each $i \in \{1, 2\}$, $K_i = J[V(C_i)]$ is a clique of J . For each $i \in \{1, 2\}$, let K'_i be a maximal clique of J containing K_i . Then we have $c_1 \in K'_1$ and $c_i \in K'_2$, which implies that K'_1 and K'_2 are distinct. But now the edge $pq \in E(J)$ is contained in two maximal cliques of J , namely K'_1 and K'_2 , which violates J being a bloated tree. This proves Lemma 5.3. ■

The following was proved in [11]:

Theorem 5.4 (Davies [11]). *For every integer $k \geq 1$, there exists an integer $f = f(k)$ such that if G is a connected graph and $S \subseteq V(G)$ with $|S| \geq f(k)$, then G has an induced subgraph J which is a bloated tree and $|J \cap S| \geq k$.*

We also need the following well-known result; see, for example, [2] for a proof.

Lemma 5.5. *For all positive integers d, q , there exists a positive integer $N(d, q)$ such that for every connected graph G on at least $N(d, q)$ vertices, either G contains a vertex of degree at least d , or there is a path in G with q vertices.*

For a graph G and a set $S \subseteq G$, by an S -bump we mean a vertex $v \in G \setminus S$ of degree two in G , say $N_G(v) = \{v_1, v_2\}$, such that $v_1 v_2 \notin E(G)$. Also, by *suppressing* the S -bump v we mean removing v from G and adding the edge $v_1 v_2$ (hence, G is a subdivision of the resulting graph). We are now ready to prove Theorem 5.2, which we restate:

Theorem 5.2. *For every integer $\eta \geq 1$, there exists an integer $\mu = \mu(\eta) \geq 1$ with the following property. Let G be a graph and $S \subseteq V(G)$ with $|S| \geq \mu$ such that S is contained in a connected component of G . Then G contains an (S, η) -connectifier H . In particular, H is connected, $|H \cap S| = \eta$, and every vertex in $H \cap S$ has degree at most η in H .*

Proof. Let $f(\cdot)$ be as in Theorem 5.4, and $N(\cdot, \cdot)$ be as in Lemma 5.5. We choose

$$\mu = \mu(\eta) = f(\max\{N(\eta, 8\eta^2 + \eta), 2\}).$$

By Theorem 5.4, since $|S| \geq \mu$, it follows that G has an induced subgraph J which is a bloated tree with $|J \cap S| \geq \max\{N(\eta, 8\eta^2 + \eta), 2\}$, and subject to this property, J has as few vertices as possible. Assume that $\eta = 2$. Then, since J is connected and $|J \cap S| \geq 2$, there is a path H in J with ends in S and $H^* \cap S = \emptyset$. But then H is an $(S, 2)$ -connectifier of type 2 in G , as desired. Therefore, we may assume that $\eta \geq 3$.

(3) *Let $X \subseteq J$ such that X is connected. Then for every connected component Q of $J \setminus X$, we have $Q \cap S \neq \emptyset$. In particular, we have $\mathcal{Z}(J) \subseteq S$.*

Suppose not. Let Q be a component of $J \setminus X$ such that $Q \cap S = \emptyset$. Since X connected, it holds that $J \setminus Q$ is connected, as well. It follows that $J \setminus Q$ is bloated tree and $|(J \setminus Q) \cap S| = |J \cap S|$, which contradicts the minimality of J . This proves (3).

Let J_1 be the graph obtained from J by successively suppressing S -bumps in J until there are none. Then J_1 is also a bloated tree, and J is a subdivision of J_1 . The following is immediate from (3) and the definition of J_1 .

(4) J_1 has no S -bump and $J_1 \cap S = J \cap S$. Also, for every $X \subseteq J_1$ with X connected and every connected component Q of $J_1 \setminus X$, we have $Q \cap S \neq \emptyset$. In particular, we have $\mathcal{Z}(J_1) \subseteq S$.

Since J is a bloated tree and so contains no hole, it follows that J is a subdivision of J_1 with the additional property that for every edge $e \in E(J_1)$ which is contained in a big clique of J_1 , we have $e \in E(J)$ (that is, e is not subdivided while obtaining J from J_1). This, along with the fact that $J_1 \cap S = J \cap S$, implies that J contains an (S, η) -connectifier if and only if J_1 contains an (S, η) -connectifier. Therefore, in order to prove Theorem 5.2, it suffices to show that J_1 contains an (S, η) -connectifier, which we do in the rest of the proof.

(5) Let K be a maximal clique of J_1 , and for every $v \in K$, let Q_v be the connected component of $J_1 \setminus (K \setminus \{v\})$ containing v . Then for every two distinct vertices $u, v \in K$, we have $Q_u \cap Q_v = \emptyset$, and uv is the only edge of J_1 between Q_u and Q_v .

Suppose for a contradiction that there exist two distinct vertices $u, v \in K$ for which either $Q_u \cap Q_v \neq \emptyset$ or there is an edge in J_1 different from uv with one end in Q_u and one end in Q_v . It follows that $J_1[Q_u \cup Q_v] - uv$ is connected, and so there exists a path P in J_1 of length more than one from u to v with $P^* \subseteq (Q_u \cup Q_v) \setminus \{u, v\} \subseteq J_1 \setminus K$. Let $x \in P^*$. Then $C = u-P-v-u$ is a cycle in J_1 . Since J_1 is a bloated tree, by Lemma 5.3, $V(C)$ is a clique, and so x is adjacent to both u and v . Now, suppose that there exists a vertex $y \in K \setminus N_{J_1}(x)$. Then we have $y \notin \{u, v\}$, and so $C' = x-u-y-v-x$ is a cycle in J_1 where $V(C')$ contains two non-adjacent vertices, namely x and y , which contradicts Lemma 5.3 and the fact that J_1 is a bloated tree. Therefore, x is complete to K , and so $K \cup \{x\}$ is a clique of J_1 strictly containing K . This violates the maximality of K , and so proves (5).

(6) Suppose that J_1 contains a big clique K with $|K| \geq \eta$. Then J_1 contains an (S, η) -connectifier of type 0.

For every $v \in K$, let Q_v be the connected component of $J_1 \setminus (K \setminus \{v\})$ containing v . Then by (5), for every two distinct vertices $u, v \in K$, we have $Q_u \cap Q_v = \emptyset$, and there is no edge in J_1 with one end in Q_u and one end in Q_v except for uv . Also, by (4), for every $v \in K$, we have $Q_v \cap S \neq \emptyset$. Therefore, since Q_v is connected, we can choose a path P_v in Q_v from v to a vertex $\ell_v \in S$ (possibly $v = \ell_v$) with $P_v \cap S = \{\ell_v\}$. It follows that for distinct $u, v \in K$, we have $P_u \cap P_v = \emptyset$, and there is no edge in J_1 with one end in P_u and one end in P_v except for uv . Now, let $K' \subseteq K$ with $|K'| = \eta$. Since $\eta \geq 3$, it follows that $H = J_1[\bigcup_{v \in K'} P_v]$ is a loosely (S, η) -tied line graph of a subdivided star; that is, H is an (S, η) -connectifier of type 0 in J_1 . This proves (6).

(7) Let $x \in J_1$ such that $N_{J_1}(x)$ is a stable set of J_1 , and for every $a \in N_{J_1}(x)$, let Q_a be the connected component of $J_1 \setminus x$ containing a . Then the sets $\{Q_a : a \in N_{J_1}(x)\}$ are pairwise disjoint and anticomplete to each other.

Suppose for a contradiction that there exist two distinct vertices $a, b \in N_{J_1}(x)$ for which either $Q_a \cap Q_b \neq \emptyset$, or there is an edge in J_1 with one end in Q_a and one end in Q_b . It follows that $J_1[Q_a \cup Q_b]$ is connected, and so there exists a path P in J_1 of length more than one from a to b with $P^* \subseteq Q_a \cap Q_b \setminus \{a, b\} \subseteq J_1 \setminus \{a, b, x\}$. Then $C = a-P-b-x-a$ is a cycle in J_1 where $V(C)$ contains two non-adjacent vertices, namely a and b . This contradicts Lemma 5.3 and the fact that J_1 is a bloated tree, and so proves (7).

Now we can handle the case where J_1 contains vertices of large degree.

(8) Suppose that J_1 has a vertex of degree at least η . Then J_1 contains an (S, η) -connectifier of type 0 or 1.

Since J_1 is a bloated tree, for every vertex $x \in J_1$, either $N_{J_1}(x)$ is a clique, or $N_{J_1}(x)$ is stable set, or $J_1[N_{J_1}(x)]$ has an isolated vertex y for which $N_{J_1}(x) \setminus \{y\}$ is a clique. Therefore, J_1 has a vertex of degree at least η , and it follows that either J_1 contains a big clique K with

$|K| \geq \eta$ or there exists a vertex $x \in V(J_1)$ of degree at least η in J_1 such that $N_{J_1}(x)$ is a stable set of J_1 . In the former case, (8) follows from (6). So we may assume that the latter case holds. For each $a \in N_{J_1}(x)$, let Q_a be the connected component of $J_1 \setminus x$ containing a . Then by (7), the sets $\{Q_a : a \in N_{J_1}(x)\}$ are pairwise disjoint and anticomplete to each other. Also, by (4), for every $a \in N_{J_1}(x)$, we have $Q_a \cap S \neq \emptyset$. Therefore, since Q_a is connected, we can choose a path P_a in Q_a from a to a vertex $\ell_a \in S$ (possibly $a = \ell_a$) with $P_a \cap S = \{\ell_a\}$. It follows that the paths $\{P_a : a \in N_{J_1}(x)\}$ are pairwise disjoint and anticomplete to each other. Let A be a subset of $N_{J_1}(x)$ with $|A| = \eta - 1$ if $x \in S$ and $|A| = \eta$ if $x \notin S$. Then $H = J_1[\bigcup_{a \in A} P_a]$ is a (S, η) -tied rooted subdivided star with root x such that $(H \cap S) \setminus \mathcal{L}(H) \subseteq \{x\}$; that is, H is (S, η) -connectifier in J_1 of type 1. This proves (8).

Henceforth, by (8), we may assume that J_1 has no vertex of degree at least η . Also, by (4), we have $|J_1| \geq |J_1 \cap S| \geq N(\eta, 8\eta^2 + \eta)$. As a result, by Lemma 5.5, J_1 contains a path P on $8\eta^2 + \eta$ vertices.

(9) *Suppose that there is no path in $P \setminus S$ of length 8η . Then J_1 contains an (S, η) -connectifier of type 2.*

Suppose not. Then P contains no (S, η) -tied path. Let $|P \cap S| = s$. It follows that $s < \eta$. Therefore, since there is no path in $P \setminus S$ of length 8η , we have $|P| \leq 8\eta(s + 1) + s < 8\eta^2 + \eta$, a contradiction. This proves (9).

In view of (9), we may assume that P contains a path P_1 of length 8η with $P_1 \cap S = \emptyset$, say

$$P_1 = d_0 - a_1 - b_1 - c_1 - d_1 - a_2 - b_2 - c_2 - d_2 - \cdots - a_{2\eta} - b_{2\eta} - c_{2\eta} - d_{2\eta}.$$

For each $i \in [2\eta]$, let $A_i = \{a_i, b_i, c_i\}$, let L_i be the connected component of $J_1 \setminus A_i$ containing $P_1[d_0, d_{i-1}]$, and let R_i be the connected component of $J_1 \setminus A_i$ containing $P_1[d_i, d_{2\eta}]$. We deduce:

(10) *For each $i \in [2\eta]$, L_i and R_i are distinct, and so $L_i \cap R_i = \emptyset$.*

Suppose not. Then $J_1[L_i \cup R_i]$ is connected. Therefore, there exists a path Z in J_1 from a vertex $z \in L_i$ to a vertex $z' \in R_i$ such that $Z^* \subseteq (L_i \cup R_i) \setminus P_1$. But then $C = z - P_1 - z' - Z - z$ is a cycle in J_1 and $V(C)$ contains two non-adjacent vertices, namely a_i and c_i , contradicting that J_1 is a bloated tree. This proves (10).

(11) *For each $i \in [2\eta]$, there exists a component Q_i of $J_1 \setminus A_i$ different from L_i and R_i .*

Suppose not. Then $J_1 \setminus A_i$ has exactly two distinct components, namely L_i and R_i . Assume that b_i has degree two in J_1 . Then, since $b_i \in P_1 \subseteq J_1 \setminus S$, it follows that b_i is an S -bump, which violates (4). So there exists a vertex $z \in N_{J_1}(b_i) \setminus A_i \subseteq L_i \cup R_i$, say $z \in L_i$. Consequently, since L_i is connected, there exists a path Z in L_i from z to a vertex $z' \in P_1[d_0, d_{i-1}]$ with $Z \cap P_1 = \{z'\}$. But then $C = b_i - z - Z - z' - P_1 - b_i$ is a cycle in J_1 and $V(C)$ contains two non-adjacent vertices, namely b_i and d_{i-1} , contradicting that J_1 is a bloated tree. This proves (11).

(12) *For each $i \in [2\eta]$, let Q_i be as in (11). Then we have $P_1 \cap Q_i = \emptyset$ and $N_{J_1}(Q_i) \subseteq A_i$. Also, the sets $\{Q_i : i \in [2\eta]\}$ are pairwise disjoint and anticomplete to each other.*

The first two assertions are immediate from the fact that Q_i is a component of $J_1 \setminus A_i$ different from L_i and R_i . For the third one, suppose for a contradiction that $Q_i \cup Q_j$ is connected for some distinct $i, j \in [2\eta]$, say $i < j$. Since J_1 is connected and $N_{J_1}(Q_j) \subseteq A_j$, it follows that $Q_j \cup A_j$ is connected, and so $Q_i \cup Q_j \cup A_j$ is connected. As a result, there exists a path R in J_1 with one end $q \in Q_i$ and one end $q' \in A_j \subseteq R_i$ with $R^* \subseteq Q_j$. Also, we have $A_i \cap R \subseteq A_i \cap (Q_i \cup Q_j \cup A_j) \subseteq P_1 \cap (Q_i \cap Q_j) = \emptyset$. In other words, R is a path in $J_1 \setminus A_i$ from $q \in Q_i$ to $q' \in L_i$. But then we have $q \in Q_i \cap R_i$, a contradiction with (11). This proves (12).

For each $i \in [2\eta]$, let Q_i be as in (11). Then by (4), since A_i is connected, we have $Q_i \cap S \neq \emptyset$. Also, from (11) and the connectivity of J_1 , we have $N_{Q_i}(A_i) \neq \emptyset$. Therefore, since Q_i is connected, we can choose a path W_i in Q_i from a vertex in $x_i \in N_{Q_i}(A_i)$ to a vertex in $y_i \in Q_i \cap S$

(possibly $x_i = y_i$) such that $W_i^* \cap (N_{Q_i}(A_i) \cup S) = \emptyset$. Let $G_i = J_1[A_i \cup W_i]$. It follows that G_i is connected and $G_i \cap S = \{y_i\}$.

The following is easily observed:

(13) *The sets $\{G_i : i \in [2\eta]\}$ are pairwise disjoint and anticomplete to each other. Also, for every $i \in [2\eta]$, $d_{i-1}a_i$ and $c_i d_i$ are the only edges in J_1 with one end in G_i and one end in $P_1 \setminus G_i$.*

The proof is almost concluded. Note that since J_1 is a bloated tree, it follows that for every $i \in [2\eta]$, there is no cycle in J_1 containing both a_i and c_i . Consequently, we have either $|N_{A_i}(x_i)| = 1$, or $N_{A_i}(x_i) = \{a_i, b_i\}$, or $N_{A_i}(x_i) = \{b_i, c_i\}$, as otherwise $x_i - a_i - b_i - c_i - x_i$ is a cycle in J_1 containing both a_i and c_i . Let $I \subseteq [2\eta]$. We say I is *light* if $|N_{A_i}(x_i)| = 1$ for every $i \in I$. Also, we say I is *heavy* if for every $i \in I$, we have either $N_{A_i}(x_i) = \{a_i, b_i\}$, or $N_{A_i}(x_i) = \{b_i, c_i\}$. It follows that there exists $I \subseteq [2\eta]$ with $|I| = \eta$ which is either light or heavy. Let i_1 and i_η be smallest and the largest elements of I , respectively. It follows from $\eta \geq 3$ that i_1 and i_η are distinct and $i_\eta \geq 3$. Let Z_1 be a path in G_{i_1} from c_{i_1} to y_{i_1} , and let Z_η be a path in G_{i_η} from a_{i_η} to y_{i_η} . Let

$$H = J_1 \left[P_1[c_{i_1}, a_{i_\eta}] \cup (Z_1 \cup Z_\eta) \cup \left(\bigcup_{i \in I \setminus \{i_1, i_\eta\}} G_i \right) \right].$$

Using (13), it is straightforward to observe that if I is light, then H is a loosely (S, η) -tied caterpillar, and if I is heavy, then H is a loosely (S, η) -tied line graph of a caterpillar. In other words, H is an (S, η) -connectifier of type 3 or 4. This completes the proof of Theorem 5.2. ■

6. STRONG k -BLOCKS

Let G be a graph. By a *separation* in G we mean a triple (L, M, R) of pairwise disjoint subsets of vertices in G with $L \cup M \cup R = G$, such that neither L nor R is empty and L is anticomplete to R in G . Let $x, y \in G$ be distinct. We say a set $M \subseteq G \setminus \{x, y\}$ *separates x and y* if there exists a separation (L, M, R) in G with $x \in L$ and $y \in R$. For a positive integer k , a k -*block* in G is a maximal set B of at least k vertices such that no two distinct vertices $x, y \in B$ are separated by a set $M \subseteq G \setminus \{x, y\}$ with $|M| < k$. The application of k -blocks to bounding the treewidth in hereditary graph classes is not unprecedented; see for example, [16, 21]. However, we find it best to work with a stronger notion of a k -block, which we define next.

Let k be a positive integer and let G be a graph. A *strong k -block* in G is a set B of at least k vertices in G such that for every 2-subset $\{x, y\}$ of B , there exists a collection $\mathcal{P}_{\{x, y\}}$ of at least k distinct and pairwise internally disjoint paths in G from x to y , where for every two distinct 2-subsets $\{x, y\}, \{x', y'\} \subseteq B$ and every choice of paths $P \in \mathcal{P}_{\{x, y\}}$ and $P' \in \mathcal{P}_{\{x', y'\}}$, we have $P \cap P' = \{x, y\} \cap \{x', y'\}$.

In this section, we prove that for all positive integers k and t , every t -clean graph with no strong k -block has bounded treewidth. In other words, we show that for every positive integer k , the class of all graphs with no strong k -block is clean.

To begin with, we need some definitions as well as a couple of results from the literature. For a tree T and an edge $xy \in E(T)$, we denote by $T_{x, y}$ the component of $T - xy$ containing x . Let G be a graph and (T, χ) be a tree decomposition for G . For every $S \subseteq T$, let $\chi(S) = \bigcup_{x \in S} \chi(x)$. Also, for every edge $xy \in E(T)$, we define an *adhesion* for (T, χ) as $\chi(x, y) = \chi(x) \cap \chi(y) = \chi(T_{x, y}) \cap \chi(T_{y, x})$. For every $x \in V(T)$, by the *torso at x* , denoted by $\hat{\chi}(x)$, we mean the graph obtained from the bag $\chi(x)$ by, for each $y \in N_T(x)$, adding an edge between every two non-adjacent vertices $u, v \in \chi(x, y)$. It is a well-known observation that clique cutsets do no effect the treewidth. More precisely, the following holds (a proof can be worked out easily using Lemma 5 from [8]).

Theorem 6.1 (folklore, see Lemma 5 in [8]). *Let G be a graph and let (T, χ) be a tree decomposition for G . Then the treewidth of G is at most the maximum treewidth of a torso $\hat{\chi}(x)$ taken over all $x \in V(T)$.*

Next we bring the material we need from [12] and [22]. The *fatness* of a tree decomposition (T, χ) of an n -vertex graph G is the $(n + 1)$ -tuple (a_0, \dots, a_n) , where a_i denotes the number of parts of (T, χ) of size $n - i$. If (T, χ) has lexicographically minimum fatness among all tree decompositions with all adhesions less than k , we call (T, χ) *k-atomic*. Also, a tree decomposition (T, χ) of a graph G is *tight* if for each vertex $x \in V(T)$ and every neighbor $y \in V(T)$ of x , there is a component C of $\chi(T_{y,x}) \setminus \chi(T_{x,y})$ such that every vertex in $\chi(x, y)$ has a neighbor in C . The following is proved in [22].

Lemma 6.2 (Weißbauer, Lemma 6 in [22]). *Every k -atomic tree decomposition is tight.*

Let (T, χ) be a tree decomposition for a graph G and \mathcal{S} be a set of pairwise disjoint subtrees of T . Let T' be the tree obtained from T by contracting every subtree $S \in \mathcal{S}$ into a new vertex v_S . Let $\chi' : V(T') \rightarrow 2^{V(G)}$ be defined as follows. Let $\chi'(v_S) = \chi(S)$ for every $S \in \mathcal{S}$, and let $\chi'(v) = \chi(v)$ for every $v \in V(T') \setminus \{v_S : S \in \mathcal{S}\} = V(T) \setminus (\bigcup_{S \in \mathcal{S}} S)$. One may readily observe that (T', χ') is a tree decomposition for G , which is referred to as a *contraction of (T, χ)* . The following theorem from [12] is the key ingredient in our proof of the main result of this section.¹

Theorem 6.3 (Erde and Weißbauer [12], see also [14]). *Let r be a positive integer, and let G be a graph containing no subdivision of K_r as a subgraph. Then G admits a tree decomposition (T, χ) for which the following hold.*

- (T, χ) is a contraction of a k -atomic tree decomposition for G with $k = r(r - 1)$.
- Every adhesion of (T, χ) has cardinality less than r^2 .
- For every $x \in V(T)$, either $\hat{\chi}(x)$ has fewer than r^2 vertices of degree at least $2r^4$, or $\hat{\chi}(x)$ has no minor isomorphic to K_{2r^2} .

It is straightforward to check that every contraction of a tight tree decomposition is tight. Also, for every positive integer k and every graph G , if G contains a subdivision of K_{k^3} as a subgraph, then G contains a strong k -block. Therefore, the following is immediate from Theorem 6.3 and Lemma 6.2.

Theorem 6.4. *Let k be a positive integer and let G be a graph containing no strong k -block. Then G admits a tight tree decomposition (T, χ) for which the following hold.*

- Every adhesion of (T, χ) has cardinality less than k^6 .
- For every $x \in V(T)$, either $\hat{\chi}(x)$ has fewer than k^6 vertices of degree at least $2k^{12}$, or $\hat{\chi}(x)$ has no minor isomorphic to K_{2k^6} .

We can now prove the main result of this section. For every positive integer k , let \mathcal{B}_k be the class of all graphs with no strong k -block.

Theorem 6.5. *For every integer $k \geq 1$, the class \mathcal{B}_k is clean.*

Proof. Let $t \geq 1$ and let $G \in \mathcal{B}_k^t$, that is, G is a t -clean graph with no strong k -block. We aim to show that there exists an integer $w(k, t) \geq 1$ such that $\text{tw}(G) \leq w(k, t)$. By Theorem 6.4, G has a tight tree decomposition (T, χ) for which every torso either has fewer than k^6 vertices of degree at least $2k^{12}$ or has no minor isomorphic to K_{2k^6} . For each $x \in V(T)$, let $K_x \subseteq \hat{\chi}(x)$ be the set of all vertices in $\hat{\chi}(x)$ of degree at least $2k^{12}$. We define τ_x as follows: if $|K_x| < k^6$, then let $\tau_x = \hat{\chi}(x) \setminus K_x$, and otherwise let $\tau_x = \hat{\chi}(x)$. It follows that either τ_x has maximum degree less than $2k^{12}$, or τ_x has no minor isomorphic to K_{2k^6} . Let $\xi(\cdot, \cdot)$ be as in Theorem 1.2 and $\gamma(\cdot, \cdot)$ be as in Theorem 1.3. Let

$$\begin{aligned}\gamma_0 &= \gamma(3, t), \\ \gamma_1 &= \gamma(2k^{12}, 2\gamma_0),\end{aligned}$$

¹We remark that the corresponding statement in [12], namely ‘‘Theorem 4’’ therein, does not explicitly mention that (T, χ) is a contraction of a k -atomic tree decomposition. However, as the reader can check, the proof given in Section 3 of [12] is easily seen to yield this: it starts with a k -atomic tree decomposition ‘‘ (T, \mathcal{V}) ’’ with $k = r(r - 1)$, and concludes at the end that the desired tree decomposition is a certain contraction of (T, \mathcal{V}) .

$$\begin{aligned}\xi_1 &= \xi(K_{2k^6}, 2\gamma_0), \\ w_1 &= w_1(k, t) = \max\{\gamma_1, \xi_1\}.\end{aligned}$$

We claim that:

$$(14) \text{ For every } x \in V(T), \text{ we have } \text{tw}(\tau_x) \leq w_1.$$

Suppose for a contradiction that $\text{tw}(\tau_x) > w_1$ for some $x \in V(T)$. Note that either τ_x has maximum degree less than $2k^{12}$ or τ_x has no minor isomorphic to K_{2k^6} . Therefore, the choice of w_1 together with Theorems 1.2 and 1.3 implies that τ_x contains an induced subgraph W which is isomorphic to either a subdivision of $W_{2\gamma_0 \times 2\gamma_0}$ or the line graph of a subdivision of $W_{2\gamma_0 \times 2\gamma_0}$. On the other hand, it can be seen that for every positive integer q , every subdivision of $W_{2q \times 2q}$ contains an induced subgraph isomorphic to a proper subdivision of $W_{q \times q}$ (see Figure 5.) Consequently, W , and so τ_x , contains an induced subgraph W_0 which is isomorphic to either a proper subdivision of $W_{\gamma_0 \times \gamma_0}$ or the line graph of a proper subdivision of $W_{\gamma_0 \times \gamma_0}$. In particular, W_0 has maximum degree at most three. Let us say a non-empty subset $K \subseteq W_0$ is a *blossom* if there exists $y \in N_T(x)$ such that $K \subseteq \chi(x, y)$, and subject to this property, K is maximal with respect to inclusion. It follows that every blossom K is a clique in W_0 and so we have $|K| \in \{1, 2, 3\}$. Also, every two blossoms intersect in at most one vertex, and since no two triangles in W_0 share a vertex, blossoms of cardinality three are pairwise disjoint. Let \mathcal{K} be the set of all blossoms, and for every blossom $K \in \mathcal{K}$, let us fix $y_K \in N_T(x)$ such that $K \subseteq \chi(x, y_K)$. From the maximality of blossoms, it follows that the vertices $\{y_K : K \in \mathcal{K}\}$ are all distinct. Note that (T, χ) is tight, and so for every $y \in N_T(x)$, there exists a component $C(y)$ of $\chi(T_{y,x}) \setminus \chi(T_{x,y})$ such that the every vertex in $\chi(x, y)$ has a neighbor in $C(y)$. Since (T, χ) is a tree decomposition, it follows that the sets $\{C(y_K) : y_K \in N_T(x)\}$ are pairwise distinct, disjoint and anticomplete in G . Let H_K be a connected induced subgraph of $G[(C(y_K) \cup K)]$ which contains K , and subject to this property, assume that H_K is minimal with respect to inclusion. It follows that if $|K| = 1$, then $H_K = K$, if $|K| = 2$, then H_K is a path in G between the two vertices in K with $H_K^* \subseteq C(y_K)$, and if $|K| = 3$, then H_K satisfies one of the two outcomes of Theorem 5.1. Also, the sets $\{H_K \setminus K : K \in \mathcal{K}\}$ are pairwise distinct, disjoint and anticomplete in G . Now, let

$$H = G \left[\left(W_0 \setminus \left(\bigcup_{K \in \mathcal{K}} K \right) \right) \cup \left(\bigcup_{K \in \mathcal{K}} H_K \right) \right].$$

Let H' be the minor of H obtained through the following steps in order:

- (i) For every blossom $K \in \mathcal{K}$ with $|K| = 3$, contract the connected induced subgraph H_K of H into a vertex.
- (ii) For every blossom $K \in \mathcal{K}$ with $|K| = 2$ such that K is contained in a triangle of W_0 , contract the path H_K in H into an edge between the two vertices in K .
- (iii) Contract each triangle of the resulting graph after (ii) into a vertex.

Since W_0 is isomorphic to either a proper subdivision of $W_{\gamma_0 \times \gamma_0}$ or the line graph of a proper subdivision of $W_{\gamma_0 \times \gamma_0}$, it is readily observed that H' is isomorphic to a subdivision of $W_{\gamma_0 \times \gamma_0}$. It follows that H contains $W_{\gamma_0 \times \gamma_0}$ as a minor, and so we have $\text{tw}(H) \geq \gamma_0 = \gamma(3, t) + 1$. Note that since W_0 has maximum degree at most three, H has maximum degree at most three, as well. Therefore, by Theorem 1.3, H , and so G , contains either a subdivision of $W_{t \times t}$ or the line graph of a subdivision of $W_{t \times t}$ as an induced subgraph. But this violates the assumption that G is t -clean, and so proves (14).

Now, for every $x \in V(T)$, if $|K_x| < k^6$, then we have $\hat{\chi}(x) = \tau_x \cup K_x$, and otherwise we have $\hat{\chi}(x) = \tau_x$. This, along with (14), implies that $\text{tw}(\hat{\chi}(x)) \leq w_1 + k^6$ for every $x \in V(T)$. Hence, writing $w(k, t) = w_1(k, t) + k^6$, by Theorem 6.1, we have $\text{tw}(G) \leq w(k, t)$. This completes the proof of Theorem 6.5. \blacksquare

Note that for every integer $k \geq 1$, if a graph G contains a strong k -block, then G contains K_k as a topological minor, which in turn implies that G contains every k -vertex graph as a




FIGURE 5. Proof of (14): the subgraph of $W_{8 \times 8}$ induced by the filled nodes (top) is isomorphic to a proper subdivision of $W_{4 \times 4}$ (bottom). Note the correspondence between the numbered removed nodes at the top and the inner faces of the subdivided wall at the bottom.

topological minor. Therefore, the following common strengthening of Theorems 1.2 and 1.3 is in fact an immediate corollary of Theorem 6.5:

Corollary 6.6. *For every graph H , the class of all graphs with no H -topological-minor is clean.*

7. k -BLOCKS WITH DISTANT VERTICES

The main result of this section, Theorem 7.2, asserts that for every positive integer k , every graph containing a sufficiently large block contains either a subgraph that is a subdivision of a large complete graph with all paths short, or an induced subgraph which contains a k -block with its vertices pairwise far from each other. This will be of essential use in subsequent sections, and before proving it, we recall the classical result of Ramsey (see e.g. [5] for an explicit bound).

Theorem 7.1 (See [5]). *For all integers $a, b \geq 1$, there exists an integer $R = R(a, b) \geq 1$ such that every graph G on at least $R(a, b)$ vertices contains either a clique of cardinality a or a stable set of cardinality b . In particular, for all integers $t \geq 1$ and $\rho \geq R(t, t)$, every graph G containing $K_{\rho, \rho}$ as a subgraph contains either K_t or $K_{t, t}$ as an induced subgraph.*

For a graph G and a positive integer d , a d -stable set in G is a set $S \subseteq G$ such that for every two distinct vertices $u, v \in S$, there is no path of length at most d in G from u to v . Note that a d -stable set is also a d' -stable set for every $0 < d' \leq d$. Here comes the main result of this section.

Theorem 7.2. *For all integers $d, k \geq 1$ and $m \geq 2$, there exists an integer $k_0 = k_0(d, k, m) \geq 1$ with the following property. Let G be a graph and B_0 be a strong k_0 -block in G . Assume that G does not contain a $(\leq d)$ -subdivision of K_m as a subgraph. Then there exists $A \subseteq G$ with $S \subseteq B_0 \setminus A$ such that S is both a strong k -block and a d -stable set in $G \setminus A$.*

Proof. Let $R(m, k)$ be as in Theorem 7.1. We show that

$$k_0 = k_0(d, k, m) = \binom{R(m, k)}{2} (d - 1) + R(m, k)$$

satisfies Theorem 7.2. Let $X \subseteq B_0$ with $|X| = R(m, k)$. Let $g = \binom{R(m, k)}{2}$. Let e_1, \dots, e_g be an enumeration of all 2-subsets of X , and let $e_i = \{x_i, y_i\}$ for each $i \in [g]$. Let $U_0 = \emptyset$, and for every $i \in [g]$, having defined U_{i-1} , we define P_i and U_i as follows. If there exists a path P in G of length at most d from x_i to y_i with $P^* \cap (U_{i-1} \cup X) = \emptyset$, then let $P_i = P$ and $U_i = U_{i-1} \cup P_i^*$. Otherwise, let $P_i = \emptyset$ and $U_i = U_{i-1}$. It follows that for all $i, j \in [g]$ with $i < j$ and $P_i, P_j \neq \emptyset$, we have $P_i \cap P_j^* = U_i \cap P_j^* = \emptyset$ and $P_i^* \cap P_j = P_i^* \cap X = \emptyset$.

Let G_0 be the graph with $V(G_0) = X$ and for each $i \in [g]$, x_i is adjacent to y_i in G_0 if and only if $P_i \neq \emptyset$.

(15) G_0 contains no clique of cardinality m .

Suppose for a contradiction that G_0 contains a clique C of cardinality m . Then for every $i \in [g]$ with $e_i \subseteq C$, we have $P_i \neq \emptyset$. Also, for all distinct $i, j \in [g]$, we have $P_i \cap P_j^* = P_i^* \cap P_j = \emptyset$. But then $G[\bigcup_{e_i \subseteq C} P_i]$, and so G contains a $(\leq d)$ -subdivision of K_m as a subgraph, a contradiction. This proves (15).

Since $|G_0| = |X| = R(m, k)$, it follows from Theorem 7.1 and (15) that G_0 contains a stable set S of cardinality k . Let $A = U_g \cup (X \setminus S)$. Then we have $|A| \leq g(d - 1) + R(m, k) - k$. Therefore, since $S \subseteq B_0 \setminus A$ and B_0 is a strong $(g(d - 1) + R(m, k))$ -block, we deduce that S is a strong k -block in $G \setminus A$. It remains to show that S is a d -stable set in $G \setminus A$. Suppose not. Then there exists $x, y \in S$ and a path Q in $G \setminus A$ of length at most d from x to y . Thus, we may choose $i \in [g]$ such that $e_i \subseteq Q \cap S$. Therefore, assuming $P = Q[x_i, y_i]$, we have $P^* \cap S = \emptyset$. Now P is a path in $G \setminus A$ (and so in G) of length at most d from x_i to y_i with $P^* \subseteq G \setminus (A \cup S) = G \setminus (U_g \cup X) \subseteq G \setminus (U_{i-1} \cup X)$. It follows that $P_i \neq \emptyset$. But we have $e_i \subseteq S$ and S is a stable set in G_0 , which implies that $P_i = \emptyset$, a contradiction. This completes the proof of Theorem 7.2. \blacksquare

8. PLANTED SUBDIVIDED STAR FORESTS

In this section we extend ideas from [16] to produce a subdivided star forest whose roots are contained in sets with useful properties. Let G be a graph, $S \subseteq G$, and F a subdivided star forest. We say a subgraph F' of G isomorphic to F is S -planted if F' is rooted and $\mathcal{R}(F') \subseteq S$. Write \mathcal{H}_λ for the class of graphs with no holes of length greater than λ . The main result of this section is the following.

Theorem 8.1. *For all positive integers d, k, t, δ, λ , and θ with $\delta \geq 2$, there exists a positive integer $k_1 = k_1(d, k, t, \delta, \lambda, \theta)$ with the following property. Let G be a t -clean graph and let B_1 be a strong k_1 -block in G . Then there exist $A \subseteq V(G)$ and $S \subseteq B_1 \setminus A$ such that the following hold.*

- S is both a strong k -block and a d -stable set in $G \setminus A$.
- $G \setminus A$ contains an S -planted copy of $\theta S_{\delta, \lambda}$.
- $G \setminus A$ contains a hole of length greater than λ .

In particular, we have $\mathcal{F}_{\theta S_{\delta, \lambda}}^t, \mathcal{H}_\lambda^t \subseteq \mathcal{B}_{k_1}$.

Note that Theorem 8.1, combined with Theorem 6.5 and Lemma 2.1, implies Theorems 1.4 and 1.5 at once. Theorem 8.1 is also a key tool in the proof of Theorem 4.1 in Section 9. We need the following two results from [16].

Lemma 8.2 (Lozin and Razgon [16]). *For all positive integers a and b , there is a positive integer $c = c(a, b)$ such that if a graph G contains a collection of c pairwise disjoint subsets of $V(G)$, each of cardinality at most a and with at least one edge between every two of them, then G contains $K_{b, b}$ as a subgraph.*

Theorem 8.3 (Lozin and Razgon [16]). *For all positive integers p and r , there exists a positive integer $m = m(p, r)$ such that every graph G containing a $(\leq p)$ -subdivision of K_m as a subgraph contains either $K_{p,p}$ as a subgraph or a proper $(\leq p)$ -subdivision of $K_{r,r}$ as an induced subgraph.*

We deduce the following lemma.

Lemma 8.4. *For every integer $t \geq 1$, there exists an integer $n = n(t) \geq 1$ with the following property. Let G be a t -clean graph and let ρ be an integer with $\rho \geq R(t, t)$, where $R(\cdot, \cdot)$ is as in Theorem 7.1. Then G does not contain a $(\leq \rho)$ -subdivision of K_n as a subgraph.*

Proof. Let $n = n(t) = m(R(t, t), 2t^2)$, where $m(\cdot, \cdot)$ is as in Theorem 8.3. Suppose for a contradiction that G contains a $(\leq \rho)$ -subdivision of K_n as a subgraph. Then by Theorem 8.3, G either contains $K_{\rho, \rho}$ as a subgraph, or contains an induced subgraph H isomorphic to a proper subdivision of $K_{2t^2, 2t^2}$. In the former case, by Theorem 7.1, G contains either K_t or $K_{t,t}$, which violates the assumption that G is t -clean. In the latter case, note that a proper subdivision of $K_{2t^2, 2t^2}$ contains a proper subdivision of every bipartite graph on at most $2t^2$ vertices. In particular, H , and so G , contains a subdivision of $W_{t \times t}$, again contradicting that G is t -clean. This proves the Lemma 8.4. \blacksquare

We are now ready to prove the main result of this section.

Proof of Theorem 8.1. Let $R(\cdot, \cdot)$ be as in Theorem 7.1. Let $c = c(\lambda, R(t, t))$, where $c(\cdot, \cdot)$ is as in Lemma 8.2. Let $n = n(t)$, be as in Lemma 8.4. Let $k_0(\cdot, \cdot, \cdot)$ be as in Theorem 7.2. Let

$$k_1 = k_1(d, k, t, \delta, \lambda, \theta) = k_0(\max\{d, R(t, t), 2\lambda + 1\}, \max\{k, R(c, \delta), \theta\}, n).$$

We claim that this choice of k_1 satisfies Theorem 8.1. To see this, suppose that G is a t -clean graph which has a strong k_1 -block B_1 . Note first that, by Lemma 8.4, G does not contain a $(\leq \max\{d, R(t, t), 2\lambda + 1\})$ -subdivision of K_n as a subgraph. Therefore, by Theorem 7.2, there exist $A \subseteq G$ and $S \subseteq B_1 \setminus A$ such that S is both a strong $\max\{k, R(c, \delta), \theta\}$ -block and a $\max\{d, R(t, t), 2\lambda + 1\}$ -stable set in $G \setminus A$. In particular, S is both a strong k -block and a d -stable set in $G \setminus A$, which proves the first bullet of Theorem 8.1. Next we claim that:

(16) *For every $x \in S$, there exists a copy F_x of $S_{\delta, \lambda}$ in $G \setminus A$ where $x \in F_x$ has degree δ in F_x .*

It is easily seen that $|S| \geq 2$. Pick a vertex $y \in S \setminus \{x\}$. Since S is a strong $R(c, \delta)$ -block in $G \setminus A$, there exists a collection $\{P_i : i \in [R(c, \delta)]\}$ of pairwise internally disjoint paths in $G \setminus A$ from x to y . Since S is a $(2\lambda + 1)$ -stable set in $G \setminus A$, for each $i \in [R(c, \delta)]$, P_i has length greater than $\lambda + 1$. Let P'_i be the subpath of P_i of length λ containing x as an end. Then $\{P'_i : i \in [R(c, \delta)]\}$ is a collection of $R(c, \delta)$ pairwise disjoint subsets of $G \setminus A$, each of cardinality λ . Let Γ be the graph with $V(\Gamma) = [R(c, \delta)]$ such that for all distinct $i, j \in [R(c, \delta)]$, i is adjacent to j in Γ if and only if $P'_i \setminus \{x\}$ is not anticomplete to $P'_j \setminus \{x\}$ in G . By Theorem 7.1, Γ contains either a clique of cardinality c or a stable set of cardinality δ . Suppose first that Γ contains a clique of cardinality c . Then Lemma 8.2 implies that G contains $K_{R(t, t), R(t, t)}$ as a subgraph, and thus by Theorem 7.1, G contains K_t or $K_{t, t}$, which violates the assumption that G is t -clean. Consequently, Γ has a stable set I of cardinality δ . But now $F_x = G[\bigcup_{i \in I} P'_i]$ is a copy of $S_{\delta, \lambda}$ in $G \setminus A$ where $x \in F_x$ has degree δ in F_x . This proves (16).

Now we can prove the second bullet of Theorem 8.1. For every $x \in S$, let F_x be as in (16). Note that since S is a $(2\lambda + 1)$ -stable set in $G \setminus A$, it follows that for all distinct $x, x' \in S$, F_x and $F_{x'}$ are disjoint and anticomplete to each other. Also, since S is a strong θ -block, there exists $S' \subseteq S$ with $|S'| = \theta$. But now $G[\bigcup_{x \in S'} F_x]$ is an S' -planted copy of $\theta S_{\delta, \lambda}$ in $G \setminus A$, as desired.

It remains to prove the third bullet of Theorem 8.1. Proceeding as in the proof of (16), we choose distinct vertices $x, y \in S$ and two internally disjoint paths P_1 and P_2 in $G \setminus A$ from x to y such that $P'_1 \setminus \{x\}$ is anticomplete to $P'_2 \setminus \{x\}$, where for each $i \in \{1, 2\}$, P'_i is the subpath of P_i of length λ containing x as an end. Traversing P_1 from x to y , let z be the first vertex in P_1^* with a neighbor in $P_2 \setminus \{x\}$ (this vertex exists, since the neighbor of y in P_1 is adjacent to $P_2 \setminus \{x\}$). Also, traversing P_2 from x to y , let $w \in P_2 \setminus \{x\}$ be the first neighbor of z in $P_2 \setminus \{x\}$. Note that

since P'_1 is anticomplete to P'_2 , it follows that either $z \notin P'_1$ or $w \notin P'_2$. But now $x-P_1-z-w-P_2-x$ is a hole in $G \setminus A$ of length at least $\lambda + 3$. This completes the proof of Theorem 8.1. \blacksquare

9. PROOF OF THEOREM 4.1

The last step in the proof of Theorem 4.1 is the following. Note that the condition $\delta \geq 3$ is due to the fact that there is only one choice of roots for subdivided star forests in which every component has a branch vertex, and so it is slightly more convenient to work with them.

Lemma 9.1. *For all positive integers $t, \delta, \lambda, \sigma, \theta$ with $\delta \geq 3$ and $\theta \geq 2$, there exists an integer $k_2 = k_2(t, \delta, \lambda, \sigma, \theta) \geq 1$ with the following property. Let G be a t -clean graph containing a strong k_2 -block. Then G contains a σ -connectification of $(\theta S_{\delta, \lambda}, \mathcal{R}(\theta S_{\delta, \lambda}))$. In other words, we have $\mathcal{C}_{\sigma, \theta S_{\delta, \lambda}, \mathcal{R}(\theta S_{\delta, \lambda})}^t \subseteq \mathcal{B}_{k_2}$.*

Proof. Let $\mu(\cdot)$ be as in Theorem 5.2. Let

$$\begin{aligned}\gamma_1 &= \mu(\max\{t, \sigma\theta, \theta + 1\}), \\ \gamma_2 &= \mu(\gamma_1), \\ \gamma_3 &= \gamma_2((2t\gamma_1 + \delta)\lambda + 1).\end{aligned}$$

Let $k_1(\cdot, \cdot, \cdot, \cdot, \cdot, \cdot)$ be as in Theorem 8.1. We define:

$$k_2 = k_2(t, \delta, \lambda, \sigma, \theta) = k_1\left(2\sigma - 1, \gamma_3 + R(t, t)\binom{\gamma_3}{2t}, t, 2t\gamma_1 + \delta, \lambda, \gamma_2\right).$$

Let B_2 be a strong k_2 -block in G . By Theorem 8.1, there exist $A \subseteq G$ and $S \subseteq B_2 \setminus A$ such that the following hold. Let $G_0 = G \setminus A$.

- S is both a strong $R(t, t)\binom{\gamma_3}{2t}$ -block and a $(2\sigma - 1)$ -stable set in G_0 .
- G_0 contains an S -planted copy F of $\gamma_2 S_{2t\gamma_1 + \delta, \lambda}$.

Then $|\mathcal{R}(F)| = \gamma_2$ and $|F| = \gamma_3$. For every $x \in \mathcal{R}(F)$, let F_x be the component of F with root x . Let W be the set of all vertices in $G_0 \setminus F$ with at least $2t$ neighbors in F .

$$(17) \text{ We have } |W| < R(t, t)\binom{\gamma_3}{2t}.$$

Suppose not. Let $q = R(t, t)\binom{\gamma_3}{2t}$ and let $w_1, \dots, w_q \in W$ be distinct. For every $i \in [q]$, let N_i be a set of $2t$ neighbors of w_i in F . It follows that there exist $I \subseteq [q]$ and $N \subseteq F$ such that $|I| = R(t, t)$, $|N| = 2t$ and $N_i = N$ for all $i \in I$. Note that since F is a forest, N contains a stable set N' of G_0 with $|N'| = t$. Also, since G_0 is t -clean, it does not contain a clique of cardinality t . Thus, by Lemma 7.1, $G_0[\{w_i : i \in I\}]$ contains a stable set N'' of cardinality t . But then $G_0[N' \cup N'']$ is isomorphic to $K_{t,t}$, which contradicts that G_0 is t -clean. This proves (17).

Let $G_1 = G_0 \setminus W$. Then G_1 is a t -clean induced subgraph of G . In order to prove Theorem 9.1, it suffices to show that G_1 contains a σ -connectification of $\theta S_{\delta, \lambda}$, which we do in the rest of the proof.

Recall that S is both a strong $(\gamma_3 + R(t, t)\binom{\gamma_3}{2t})$ -block and a $(2\sigma - 1)$ -stable set in G_0 . Thus, since $S \setminus W \subseteq G_1$, by (17), $S \setminus W$ is both a strong γ_3 -block and a $(2\sigma - 1)$ -stable set in G_1 . Also, we have $\mathcal{R}(F) \subseteq S \setminus W$. It follows that $\mathcal{R}(F)$ is a $(2\sigma - 1)$ -stable set in G_1 , and for every two distinct vertices $x, x' \in \mathcal{R}(F)$, since $|F \setminus \mathcal{R}(F)| < \gamma_3$, there is a path in $G_1 \setminus (F \setminus \mathcal{R}(F))$ from x to x' . Consequently, $G_1 \setminus (F \cup \mathcal{R}(F))$ has a component containing $\mathcal{R}(F)$. Let G_2 be the graph obtained from G_1 by contracting F_x into x for each $x \in \mathcal{R}(F)$. Then G_2 contains $G_1 \setminus (F \cup \mathcal{R}(F))$ as a spanning subgraph, and so G_2 has a component containing $\mathcal{R}(F)$. Since $\mathcal{R}(F) \geq \gamma_2 = \mu(\gamma_1)$, from Theorem 5.2 applied to G_2 and $\mathcal{R}(F)$, it follows that G_2 contains a

connected induced subgraph H_2 such that, assuming $S' = H_2 \cap \mathcal{R}(F)$, we have $|S'| = \gamma_1$ and every vertex in S' has degree at most γ_1 in H_2 . Let

$$H_1 = G_1 \left[H_2 \cup \left(\bigcup_{x \in S'} F_x \right) \right].$$

In other words, H_1 is the induced subgraph of G_1 obtained from H_2 by undoing the contraction of F_x into x for each $x \in H_2 \cap \mathcal{R}(F)$. It follows that H_1 is a connected induced subgraph of G_1 and $H_1 \cap \mathcal{R}(F) = H_2 \cap \mathcal{R}(F) = S'$. Moreover, since $\mathcal{R}(F)$ is a $(2\sigma - 1)$ -stable set in G_1 , S' is also a $(2\sigma - 1)$ -stable set in H_1 .

(18) For every $x \in S'$, we have $|N_{F_x}(H_1 \setminus F_x)| < 2t\gamma_1$.

Note that $N_{H_1 \setminus F_x}(F_x) = N_{H_2}(x)$, and so $|N_{H_1 \setminus F_x}(F_x)| \leq \gamma_1$. Also, since H_1 is an induced subgraph of G_1 , by the definition of W , no vertex in $N_{H_1 \setminus F_x}(F_x) \subseteq G_1 \setminus F$ has at least $2t$ neighbors in F_x . Therefore, we have $|N_{F_x}(H_1 \setminus F_x)| < 2t\gamma_1$. This proves (18).

The following is immediate from (18) and the fact that for every $x \in S'$, F_x is isomorphic to $S_{2t\gamma_1 + \delta, \lambda}$.

(19) For every $x \in S'$, F_x contains an induced copy F'_x of $S_{\delta, \lambda}$ containing x such that $F'_x \setminus \{x\}$ is anticomplete to $H_1 \setminus F'_x$.

Next, we define:

$$H'_1 = H_1 \setminus \left(\bigcup_{x \in S'} (F'_x \setminus \{x\}) \right).$$

It follows that H'_1 is a connected induced subgraph of G_1 and $S' \subseteq H'_1$ is a $(2\sigma - 1)$ -stable set in H'_1 .

(20) H'_1 , and so G_1 , contains an (S', θ, σ) -connectifier H of type i for some $i \in [4]$.

Since $|S'| \geq \gamma_1 = \mu(\max\{t, \theta\sigma, \theta + 1\})$, we can apply Theorem 5.2 to H'_1 and S' . It follows that H'_1 contains an $(S', \max\{t, \theta\sigma, \theta + 1\})$ -connectifier H' . Since S' is a $(2\sigma - 1)$ -stable set in H'_1 , $H' \cap S'$ is also a $(2\sigma - 1)$ -stable set in H' . It is straightforward to observe that if H' is of type i for $i \in \{2, 3, 4\}$, then H' , and so H'_1 , contains an (S', θ, σ) -connectifier H . Also, if H' is of type 0, then H' contains a clique of cardinality t , which violates that G_1 is t -clean. It remains to consider the case where H' is of type 1. Then H' contains an $(S', \theta + 1)$ -tied rooted subdivided star H'' with root r in which every stem has length at least σ and $(H'' \cap S') \setminus \mathcal{L}(H'') \subseteq \{r\}$. Since $\theta \geq 2$, it follows that H'' has at least three vertices and r is not a leaf of H'' . If H'' is a path with ends $h_1, h_2 \in S'$, then $\theta = 2$ and $r \in S'$. This, along with the fact that $H'' \cap S'$ is a $(2\sigma - 1)$ -stable set in H'' , implies that $H = H''[h_1, r]$ has length at least 2σ . But then H is a (S', θ, σ) -connectifier of type 2 in H'' , and so in H'_1 . Also, if H'' is not a path, then r is the unique branch vertex of H'' . Again, since $H'' \cap S'$ is $(2\sigma - 1)$ -stable set in H' (and so in H''), there exists a stem P of H'' such that every stem of H'' other than P has length at least σ . Therefore, $H = H' \setminus (P \setminus \{r\})$ is an (S', θ, σ) -connectifier of type 1 in H'' , and so in H'_1 . This proves (20).

Let H be as in (20). Let $X = H \cap S$. Let $F' = \bigcup_{x \in X} F'_x$ and $\Xi = G_1[H \cup F']$. Then by (19), F' is an induced subgraph of Ξ isomorphic to $\theta S_{\delta, \lambda}$ and $F' \setminus X$ is anticomplete to $\Xi \setminus F'$. Also, we have $\Xi \setminus (F' \setminus X) = H$. But then by (20), Ξ is a σ -connectification of (F', X) , and so Ξ is an induced subgraph of G isomorphic to a σ -connectification of $(\theta S_{\delta, \lambda}, \mathcal{R}(\theta S_{\delta, \lambda}))$. This completes the proof of Lemma 9.1. \blacksquare

We need one more definition before proving Theorem 4.1. For two rooted subdivided star forests F_1 and F_2 , we say F_2 embeds in F_1 if $\mathcal{R}(F_2) \subseteq \mathcal{R}(F_1)$ and there exists a collection \mathcal{S} of stems of F_1 such that $F_2 = F_1 \setminus ((\bigcup_{P \in \mathcal{S}} P) \setminus \mathcal{R}(F_1))$.

Now we prove Theorem 4.1, which we restate:

Theorem 4.1. *Let $\sigma \geq 1$ be an integer, let F be a rooted subdivided star forest of size at least two and let $\pi : [|\mathcal{R}(F)|] \rightarrow \mathcal{R}(F)$ be a bijection. Then the class $\mathcal{C}_{\sigma, F, \mathcal{R}(F), \pi}$ is clean.*

Proof. Let F be of maximum degree $\delta \geq 0$, reach $\lambda \geq 0$ and size $\theta \geq 2$. For every $x \in \mathcal{R}(F)$, let F_x be the component of F with root x . Let $F^+ = \theta S_{\delta+3, \lambda+1}$ be rooted (with its unique choice of roots). For every $y \in \mathcal{R}(F^+)$, let F_y^+ be the component of F^+ with root y . Then for every $x \in \mathcal{R}(F)$ and every $y \in \mathcal{R}(F^+)$, F_y^+ contains a copy $F_{x,y}^+$ of F_x such that $F_{x,y}^+$ embeds in F_y^+ . Now, for every choice of bijections $\pi : [\theta] \rightarrow \mathcal{R}(F)$ and $\pi^+ : [\theta] \rightarrow \mathcal{R}(F^+)$, and every σ -connectification Ξ^+ of $(F^+, \mathcal{R}(F^+))$ with respect to π^+ , let

$$\Xi = (\Xi^+ \setminus F^+) \cup \left(\bigcup_{i \in [\theta]} F_{\pi(i), \pi^+(i)}^+ \right).$$

It follows that Ξ is isomorphic to a σ -connectification of $(F, \mathcal{R}(F))$ with respect to π . In other words, for every bijection $\pi : [\theta] \rightarrow \mathcal{R}(F)$, every σ -connectification of $(F^+, \mathcal{R}(F^+))$ contains an induced subgraph isomorphic to a σ -connectification of $(F, \mathcal{R}(F))$ with respect to π . Therefore, we have $\mathcal{C}_{\sigma, F, \mathcal{R}(F), \pi} \subseteq \mathcal{C}_{\sigma, F^+, \mathcal{R}(F^+)}$. This, together with Lemma 9.1, implies that for every integer $t \geq 1$, we have $\mathcal{C}_{\sigma, F, \mathcal{R}(F), \pi}^t \subseteq \mathcal{C}_{\sigma, F^+, \mathcal{R}(F^+)}^t \subseteq \mathcal{B}_{k_2}$, where $k_2 = k_2(t, \delta + 3, \lambda + 1, \sigma, \theta)$ is as in Lemma 9.1. Now the result follows from Theorem 6.5 and Lemma 2.1. \blacksquare

10. ACKNOWLEDGEMENT

We thank Rose McCarty for bringing Theorem 6.3 to our attention, and Daniel Weißauer for helpful comments.

REFERENCES

- [1] P. Aboulker, I. Adler, E. J. Kim, N. L. D. Sintiari, and N. Trotignon. “On the treewidth of even-hole-free graphs.” *European Journal of Combinatorics* **98**, (2021), 103394.
- [2] T. Abrishami, M. Chudnovsky, C. Dibek, S. Hajebi, P. Rzażewski, S. Spirkl, and K. Vušković. “Induced subgraphs and tree decompositions II. Toward walls and their line graphs in graphs of bounded degree.” *arXiv:2108.01162*, (2021).
- [3] T. Abrishami, M. Chudnovsky, S. Hajebi, and S. Spirkl. “Induced subgraphs and tree decompositions IV. (Even hole, diamond, pyramid)-free graphs.” *arXiv:2203.06775*, (2022).
- [4] T. Abrishami, B. Alecu, M. Chudnovsky, S. Hajebi, and S. Spirkl. “Induced subgraphs and tree decompositions V. One neighbor in a hole” *arXiv:2205.04420*, (2022).
- [5] M. Ajtai, J. Komlós and E. Szemerédi. “A note on Ramsey numbers.” *J. Combinatorial Theory, Ser. A* **29**, (1980), 354–360.
- [6] B. Alecu, M. Chudnovsky, S. Hajebi, and S. Spirkl. “Induced subgraphs and tree decompositions XIII. Basic obstructions in \mathcal{H} -free graphs for finite \mathcal{H} ,” *manuscript* (2023).
- [7] M. Bonamy, É. Bonnet, H. Déprés, L. Esperet, C. Geniet, C. Hilaire, S. Thomassé and A. Wesolek, “Sparse graphs with bounded induced cycle packing number have logarithmic treewidth”, *arXiv:2206.00594*, (2022).
- [8] H. Bodlaender and A. Koster. “Safe separators for treewidth.” *Discrete Mathematics* **306**, 3 (2006), 337–350.
- [9] H. L. Bodlaender. Treewidth: Structure and Algorithms. *SIROCCO 2007: Proceedings of the 14th International Colloquium on Structural Information and Communication Complexity*, (2007), 11–25.
- [10] J. Davies, appeared in an Oberwolfach technical report *DOI:10.4171/OWR/2022/1*.
- [11] J. Davies, “Vertex-minor-closed classes are χ -bounded.” *arXiv:2008.05069*, (2020).
- [12] J. Erde and D. Weißauer. “A short derivation of the structure theorem for graphs with excluded topological minors.” *SIAM Journal of Discrete Mathematics* **33**, 3 (2019), 1654–1661.
- [13] P. Gartland, D. Lokshantov, M. Pilipczuk, M. Pilipczuk, P. Rzażewski. “Finding large induced sparse subgraphs in $C_{>t}$ -free graphs in quasipolynomial time,” *STOC 2021: Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, (2021), 330–341.
- [14] M. Grohe and D. Marx. “Structure theorem and isomorphism test for graphs with excluded topological subgraphs,” *SIAM Journal on Computing* **44**, 1 (2015), 114–159.
- [15] T. Korhonen, “Grid induced minor theorem for graphs of small degree.” *J. Combin. Theory Ser. B* **160** (2023), 206 – 214.
- [16] V. Lozin, I. Razgon. “Tree-width dichotomy.” *European Journal of Combinatorics* **103**, (2022), 103517.

- [17] K. Menger, “Zur allgemeinen Kurventheorie.” *Fund. Math.* **10**, 1927, 96–115.
- [18] N. Robertson and P. Seymour. “Graph minors. II. Algorithmic aspects of tree-width.” *J. of Algorithms*, **7** (3) (1986), 309–322.
- [19] N. Robertson and P. Seymour. “Graph minors. V. Excluding a planar graph.” *J. Combin. Theory Ser. B*, **41** (1) (1986), 92–114.
- [20] N.L.D. Sintiari and N. Trotignon. “(Theta, triangle)-free and (even-hole, K_4)-free graphs. Part 1: Layered wheels,” *J. Graph Theory* **97** (4) (2021), 475–509.
- [21] D. Weißauer. “In absence of long chordless cycles, large tree-width becomes a local phenomenon.” *J. Combin. Theory Ser. B* **139**, (2019) 342–352.
- [22] D. Weißauer. “On the block number of graphs.” *SIAM Journal of Discrete Mathematics* **33**, (2019), 346–357.